



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

total nitrogen in eggs after first cleavage with that at the early blastula stage. Considerable material has been collected for further work along this line.

L. F. SHACKELL

BEAUFORT, N. C.,

September 9, 1911

OBSERVATIONS ON THE INHERITANCE OF CHARACTERS IN *ZEA MAYS* LINN.¹

IN "Red Cuzco" and some other breeds of red maize, the red coloring matter is confined to the pericarp; being therefore a fruit character and not a seed character, it does not appear in the ear immediately resulting from a direct cross between a white female and a red male.

In a red dent breed under investigation, the red pigment occurs in the aleurone layer, and not in the pericarp. Being a seed-character, it is transmitted directly by the pollen grain to the ovule of a white breed. It behaves as a dominant to whiteness; where it meets yellowness in the same grain, it is more conspicuous than yellow. The writer has not met with a previous record of the occurrence of a red pigment of this character in the aleurone layer of the maize grain.

When this red dent is crossed with a white sugar breed the segregation, in the second generation, of the two pairs of characters redness *vs.* whiteness and starchiness *vs.* sugariness, is in approximately the following proportions:

Red	{	starchy	56.25	}	= 75%
		sugary	18.75		
White	{	starchy	18.75	}	= 25%
		sugary	6.25		

In other words:

				Red Grains	
Starchy grains	75%	of	75%	= 56.25%
Sugary grains	25%	of	75%	= 18.75%
					75.00%
				White Grains	
Starchy grains	75%	of	25%	= 18.75%
Sugary grains	25%	of	25%	= 6.25%
					25.00%

¹ Fuller details will shortly appear in the *Transactions of the Royal Society of South Africa*.

A single grain has been found on the ear studied, which distinctly shows the starchy character in one half and the sugar character in the other, a very unusual feature.

A study of row-numbers in maize-ears indicates that within certain limits the number of rows of grain on an ear is subject to fluctuating variation, which may perhaps be affected by season or food supply, or both. In more than thirty plants of Arcadia sugar-maize studied this year, each of which produced two ears on one stalk, the uppermost ear has had a different number of rows from that of the lower ear. On thirteen plants the largest number of rows occurred on the lower ear, while on eight plants the largest number was on the upper ear. In twelve plants of two ears the row-numbers were the same on both; in one case there were four more rows on one ear than on the other. Several plants of Hickory King, bearing two ears, have also produced different numbers of rows on the two ears.

The range of variation appears to be limited, however. A normally 8-rowed type ranges between 4 and 14 rows, while a normally 18-rowed type ranges between 12 and 24 rows.

The result of crossing an 8-row with an 18-row type of maize is to produce an intermediate type in the first generation, both 8-row and 18-row types practically disappear in the heterozygous form. The intermediate type bears mostly 10, 12 or 14 rows, the 12-row type greatly predominating. The experiment will be continued next year, to determine the proportion of the 8-row and 18-row types which reappear. The ears produced by the cross and the reciprocal cross are indistinguishable.

A white-cobbed breed crossed with a red-cobbed produces a red cob in the first filial generation, and so does the reciprocal cross.

J. BURTT DAVY

DEPARTMENT OF AGRICULTURE,
PRETORIA